

AD-A200 300

DTIC FILE COPY

OFFICE OF NAVAL RESEARCH

Contract N00014-83-K-0470-P00003

R&T Code NR 33359-718

Technical Report No. 92

An Infrared Spectroelectrochemical Investigation of the Ion
Pairing Reactions of the Anions and Dianions of TCNE and TCNQ

by

S. B. Khoo, J. K. Foley, C. Korzeniewski, S. Pons, and C. Marcott

Prepared for publication in J. Electroanal. Chem.

Department of Chemistry
University of Utah
Salt Lake City, UT 84112

July 15, 1988

DTIC
ELECTE
NOV 14 1988
S D
CH

Reproduction in whole, or in part, is permitted for
any purpose of the United States Government

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

88 11 10 068

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release and sale. Distribution unlimited.		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) ONR Technical Report No. 92			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION University of Utah		6b. OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry Henry Eyring Building Salt Lake City, UT 84112				7b. ADDRESS (City, State, and ZIP Code)	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (if applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-83-K-0470-P00003	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program, Code 1113 800 N. Quincy Street Arlington, VA 22217				10. SOURCE OF FUNDING NUMBERS	
				PROGRAM ELEMENT NO.	PROJECT NO.
11. TITLE (Include Security Classification) An Infrared Spectroelectrochemical Investigation of the Ion Pairing Reactions of the Anions and Dianions of TCNE and TCNQ					
12. PERSONAL AUTHOR(S) S. B. Khoo, J. K. Foley, C. Korzeniewski, S. Pons, and C. Marcott					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM 9/87 TO 7/88		14. DATE OF REPORT (Year, Month, Day) July 15, 1988	
15. PAGE COUNT 24					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) infrared spectroelectrochemistry, ion radicals		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Attached.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTC USERS				21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Stanley Pons				22b. TELEPHONE (Include Area Code) (801)581-4760	
				22c. OFFICE SYMBOL	

ABSTRACT

The infrared spectra of the dianions of tetracyanoethylene (TCNE) and tetracyanoquinodimethane (TCNQ), generated electrochemically in acetonitrile, are dependent upon the supporting electrolyte. In particular, the C≡N stretching wavenumbers are higher in the presence of alkali metal salts than in the presence of tetraalkylammonium salts, and the magnitude of the shift is correlated with a positive shift in the half-wave potential of the second reduction wave. The shifts in wavenumber and half-wave potential are attributed to contact ion pairing between the dianion and the alkali metal cation. No such shifts are observed for the anion radicals of TCNE and TCNQ in acetonitrile, nor for the dianion of TCNE in dimethylsulfoxide.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
BY	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A-1	

AN INFRARED SPECTROELECTROCHEMICAL INVESTIGATION OF THE ION PAIRING REACTIONS OF THE ANIONS AND DIANIONS OF TCNE AND TCNQ

S.B. KHOO *, JOHN F. FOLEY **, CAROL KORZENIEWSKI and STANLEY PONS ***
Department of Chemistry, University of Utah, Salt Lake City, UT 84112 (U.S.A.)

CURTIS MARCOTT
Proctor and Gamble Corporation, Cincinnati, OH (U.S.A.)
(Received 8th December 1986; in revised form 13th March 1987)

ABSTRACT

(C-N triple bond)
The infrared spectra of the dianions of tetracyanoethylene (TCNE) and tetracyanoquinodimethane (TCNQ), generated electrochemically in acetonitrile, are dependent upon the supporting electrolyte. In particular, the $\text{C}\equiv\text{N}$ stretching wavenumbers are higher in the presence of alkali metal salts than in the presence of tetraalkylammonium salts, and the magnitude of the shift is correlated with a positive shift in the half-wave potential of the second reduction wave. The shifts in wavenumber and half-wave potential are attributed to contact ion pairing between the dianion and the alkali metal cation. No such shifts are observed for the anion radicals of TCNE and TCNQ in acetonitrile, nor for the dianion of TCNE in dimethylsulfoxide.


INTRODUCTION

This paper reports infrared spectra of the anion radicals and dianions of tetracyanoquinodimethane (TCNQ) and tetracyanoethylene (TCNE) in acetonitrile (AN) and dimethylsulphoxide (DMSO) solutions. The anions and dianions were generated electrochemically and the spectra were obtained by the SNIPTIRS method [1].

* Present address: Department of Chemistry, National University of Singapore, Kent Ridge, Singapore.

** Present address: Department of Chemistry, Box 4260, Texas Tech University, Lubbock, TX 79409-4260, U.S.A.

*** To whom correspondence should be addressed.



The structures of TCNQ and TCNE are shown ^{above} below. In aprotic solvents the electrochemical reduction of TCNQ and TCNE takes place in two one-electron waves, forming a stable anion radical in the first, and a stable dianion in the second.



A homogeneous recombination reaction can also take place between the dianion and the neutral molecule to give two radical anions:



In these experiments, however, this layer of solution ^{was} thin enough that all species within it were at equilibrium with the electrode, and since equilibrium (3) lies well to the right it could be ignored. Experiments using a thicker solution layer will be described elsewhere [2].

Cyclic voltammetric experiments in this laboratory [3] have provided evidence that strong ion pairing takes place between alkali metal cations and the dianions of TCNQ and TCNE in acetonitrile solution; the half wave potentials of the second reduction waves of TCNQ and TCNE were less negative when the supporting electrolyte was 0.1 M lithium perchlorate or sodium perchlorate than when the electrolyte was 0.1 M tetraethylammonium perchlorate (TEAP) or tetra-n-butylammonium fluoroborate (TBAF), indicating that the alkali metal cations interact with, and stabilize, the dianions. The half wave potentials of the first reduction waves were independent of supporting electrolyte, indicating that the radical anions do not form ion pairs with alkali metal cations in acetonitrile.

IR spectra can differ from one environment to another, and in particular from solid to solution. The geometry of the radical ion may be distorted in the solid state, and there is also the possibility of activation of usually infrared inactive modes through a vibronic interaction mechanism, involving distortion of the highly polarizable charge cloud of the anion by the highly polarizing counter cation to produce a dipole oscillating between anion and cation. Such a mechanism has been suggested by Devlin and co-workers to explain the appearance of a strong band at about 1370 cm^{-1} in the solid state spectra of $M^+ \text{TCNE}^{2-}$, where M^+ is an alkali metal cation [4-7]. It was proposed that this band is the totally symmetric C=C stretching mode of the electron cloud perpendicular to the molecular plane. Indeed *all* of the detectable infrared bands in the 600 cm^{-1} to 4000 cm^{-1} range were attributed to totally symmetric stretching modes (A_g or D_{2h}) activated by this charge oscillation mechanism. The situation is further complicated by the strong possibility that the TCNE anion radical exists as a monomer in solution at room temperature but as a

dimer in the solid state [8-10]. Some bands in the solid stated spectra of $(\text{Na}^+)_2\text{TCNE}^{2-}$ [7] and $\text{Na}^+\text{TCNQ}^{\cdot-}$ [11] have also been attributed to vibronic activation of totally symmetric modes.

In solution the counter cation might affect the spectrum through ion pairing. For example, Eargle and Emrich [12] have found that in the fairly nonpolar solvent dimethoxyethane the C-O stretching wavenumber of the benzophenone radical anion depends upon the nature of the counteraction, the wavenumber varying from 1568 cm^{-1} in the presence of K^+ to 1575 cm^{-1} in the presence of Li^+ . These shifts were attributed to ion pairing between the anions and cations.

Electrochemical generation of the radical ions allows high dielectric constant solvents and large organic counter ions to be used. Under which conditions ion pairs are less likely to form. Resonance Raman spectra of electrochemically generated $\text{TCNE}^{\cdot-}$ [13], $\text{TCNQ}^{\cdot-}$ [14] and TCNQ^{2-} [15] have been obtained in acetonitrile, using TBAF as supporting electrolyte. Infrared spectra of electrochemically generated $\text{TCNE}^{\cdot-}$ [16] and TCNE^{2-} [17] have been obtained by SNIPTIRS. In the case of $\text{TCNE}^{\cdot-}$ the electrode material was platinum, and the electrolytes were TBAF and LiClO_4 . Differences between the spectra in the two electrolytes were attributed to absorption of the anion only in the presence of TBAF. In the case of TCNE^{2-} the electrode material was carbon, the electrolyte was TBAF, and the spectrum was attributed to TCNE^{2-} adsorbed and in solution.

The spectra described here were obtained with LiClO_4 , NaClO_4 , TEAP, and TBAF as electrolytes, in order to study the effects of ion pairing.

EXPERIMENTAL

For most experiments the spectrometer was a Digilab Qualimatic instrument with a cooled (77K) HgCdTe detector. The spectrometer was purged with nitrogen and the cell was situated outside it. The cell and other instrumentation have been described in detail elsewhere [1]. The working electrode was a 7 mm diameter platinum, gold, or vitreous carbon disk mounted on the end of a brass shaft inside a 9 mm diameter Kel-F tube. The electrode was pushed up against a CaF_2 infrared-transparent window, forming a thin-layer cell. Thin layer voltammetry showed the average window to electrode distance to be $50\text{ }\mu\text{m}$. The window had a trapezoidal cross-section, allowing equal transmission of s- and p-polarized light. For acetonitrile solutions the reference electrode was a silver wire in contact with AgNO_3 (0.01 M) and TBAF (0.1 M) in acetonitrile. For DMSO solutions an SCE was used. All electrode potentials in acetonitrile are referred to the Ag/Ag^+ (0.01 M) electrode, and in DMSO to the SCE.

Sodium perchlorate (GFS) was recrystallized twice from triply distilled water. Anhydrous lithium perchlorate (GFS) was used as received. TBAF was prepared according to the method of Lund and Iverson [18] and recrystallized from methylene chloride and distilled water. TEAP (Eastman, reagent grade) was recrystallized twice from triply distilled water. Reagent grade TCNE (Eastman) was recrystallized twice from chlorobenzene. Reagent grade TCNQ (Eastman) was used as received. Acetonitrile (Burdick and Jackson UV grade, water content nominally 0.009%) and

DMSO (Burdick and Jackson, water content nominally 0.015%) were dried over Woelm neutral alumina (Supergrade I) before use.

For each system interferograms were collected at three potentials: one where the neutral species was stable, one where only the anion radical could coexist with the electrode, and one where only the dianion could coexist with the electrode. Usually 128 interferograms at 4 cm^{-1} resolutions were collected at each potential. Ratioing two of the three averaged interferograms gave, after Fourier transformation, difference spectra where bands pointing down arise from species present only at the more negative potential, and bands pointing up are due to species present only at the more positive potential.

RESULTS

The anions of TCNE

Figures 1, 3, and 5 show SNIFTIRS spectra of the TCNE (5 mM)/TBAF (0.1 M)/AN and TCNE (5 mM)/LiClO₄ (0.1 M)/AN systems in the C≡N stretching




Fig. 1. SNIFTIRS difference spectrum in C≡N stretching region for reduction of TCNE to TCNE^{•-} at a platinum electrode. Solution thickness ca 50 μm . Solutions contained TCNE (5 mM) in acetonitrile with 0.1 M LiClO₄ (top spectrum) and 0.1 M TBAF (bottom spectrum) as supporting electrolyte. Potential modulated from +0.5 V to -0.5 V (vs. Ag/0.01 M Ag⁺ reference).

Fig. 2. SNIFTIRS spectrum in C-C stretching region for reduction of TCNE to TCNE^{•-} at a platinum electrode. Same parameters as Fig. 1.




Fig. 3. SNIFTIRS difference spectrum in C≡N stretching region for reduction of TCNE to TCNE^{2-} at a platinum electrode. Solution thickness ca 50 μm . Same solutions as Fig. 1. Top spectrum: 0.1 M LiClO_4 as supporting electrolyte, potential modulated from +0.5 V to -1.2 V. Bottom spectrum: 0.1 M TBAF as supporting electrolyte, potential modulated from -0.5 V to -2.0 V.

Fig. 4. SNIFTIRS difference spectrum in C-C stretching region for reduction of TCNE to TCNE^{2-} at a platinum electrode. Same parameters as Fig. 3.

region. The only other detectable bands are in the 1100–1300 cm^{-1} region, and these are shown in Figs. 2, 4, and 6. (Noise spikes at 1160 and 1230 cm^{-1} have been subtracted out of the TBAF spectra in Figs. 4 and 6.) A platinum electrode was used to generate the anions for these spectra, but similar spectra were obtained using pyrolytic graphite and gold electrodes. All bands could be observed with either s-polarized or p-polarized radiation, so all must arise from species which are in solution rather than on the electrode surface. Bands obtained in the presence of LiClO_4 were usually weaker than bands obtained in the presence of TBAF, apparently because some precipitation of Li_2TCNE salt often takes place when TCNE^{2-} is generated in the presence of Li^+ . When TEAP was used as electrolyte the spectra were identical to those obtained with TBAF as electrolyte.

Figures 1 and 2 show SNIFTIRS spectra taken at -0.4 V (vs. Ag/Ag^+), where TCNE^{2-} is stable, with reference potential +0.4 V, where TCNE is stable. The C≡N stretching bands of TCNE are apparently too weak to be seen, but there are negative going bands from TCNE^{2-} at 2187 and 2146 cm^{-1} . At lower wavenumbers




Fig. 5. SNIFTIRS difference spectrum in C≡N stretching region for reduction of $\text{TCNE}^{\cdot-}$ to TCNE^{2-} at a platinum electrode. Solution thickness ca $50\ \mu\text{m}$. Same solutions as Fig. 1. Top spectrum: $0.1\ \text{M}$ LiClO_4 as supporting electrolyte, potential modulated from $-0.5\ \text{V}$ to $-1.2\ \text{V}$. Bottom spectrum: $0.1\ \text{M}$ TBAF as supporting electrolyte, potential modulated from $-0.5\ \text{V}$ to $-2.0\ \text{V}$.

Fig. 5. SNIFTIRS difference spectrum in C-C stretching region for reduction of $\text{TCNE}^{\cdot-}$ to TCNE^{2-} at a platinum electrode. Same parameters as Fig. 5.

there is a positive going band from TCNE at $1155\ \text{cm}^{-1}$, and a negative going band from $\text{TCNE}^{\cdot-}$ at $1180\ \text{cm}^{-1}$. The wavenumbers of all of these bands are independent of electrolyte.

Figures 3 and 4 show SNIFTIRS spectra for electrogeneration of TCNE^{2-} in the presence of TBAF and LiClO_4 . The reference potential was again $+0.4\ \text{V}$. While the spectrum of $\text{TCNE}^{\cdot-}$ in acetonitrile is independent of electrolyte, this is clearly not the case for TCNE^{2-} . With TBAF as electrolyte the dianion has two C≡N stretch bands at 2141 and $2075\ \text{cm}^{-1}$, and a band at $1207\ \text{cm}^{-1}$.

Spectra taken with reference potential $-0.4\ \text{V}$ where the anion radical is stable and with working potentials where the dianion is stable are shown in Figs. 5 and 6. As expected, positive going bands from $\text{TCNE}^{\cdot-}$ are observed at 2187 , 2146 and $1180\ \text{cm}^{-1}$ with both electrolytes, and negative going bands from TCNE^{2-} are observed at 2141 , 2075 , and $1207\ \text{cm}^{-1}$ with TBAF electrolyte and at 2175 , 2102 , and $1232\ \text{cm}^{-1}$ with LiClO_4 .

The anions of TCNQ

SNIFTIRS spectra in the C≡N stretching region for the TCNQ (5 mM)/TBAF (0.1 M)/AN and TCNQ (5 mM)/LiClO₄ (0.1 M)/AN systems are shown in Figs. 7 and 8. TEAP and NaClO₄ were also used as electrolytes; TEAP again gave the same spectra as TBAF. A platinum electrode was used for all of these experiments. All bands reported here are due to solution species, because all could be observed with either s- or p-polarized radiation.

The spectrum of TCNQ^{•-} (Fig. 7), taken with working potential at -0.2 V and reference potential +0.4 V, shows two C≡N str bands for TCNQ^{•-}, at 2182 and 2156 cm⁻¹, and a weaker band for TCNQ at 2224 cm⁻¹ (see Table 1).




Fig. 7. SNIFTIRS difference spectrum in C≡N stretching region for reduction of TCNQ to TCNQ^{•-} at a platinum electrode. Solution thickness ca 50 μm. Solutions contained TCNQ (5 mM) in acetonitrile with 0.1 M LiClO₄ (top spectrum) and 0.1 M TBAF (bottom spectrum) as supporting electrolyte. Potential modulated from +0.4 V to -0.2 V (vs. Ag/0.01 M Ag⁺ reference).

Fig. 8. SNIFTIRS difference spectrum in C≡N stretching region for reduction of TCNQ to TCNQ²⁻ at a platinum electrode. Solution thickness ca 50 μm. Same solutions as Fig. 7. Top spectrum: 0.1 M LiClO₄ as supporting electrolyte, potential modulated from +0.4 V to -0.2 V. Bottom spectrum: 0.1 M TBAF as supporting electrolyte, potential modulated from -0.4 V to -1.2 V.

TABLE 1

Infrared vibrational wavenumbers of TCNE and its anions in acetonitrile

Mode	Vibrational wavenumber/cm ⁻¹				
	TCNE ^a	TCNE ^b	TCNE ⁻ ^{b,c}	TCNE ²⁻ /TBA ⁺ ^{b,d}	TCNE ²⁻ /Li ⁺ ^b
$\nu_{15}(\text{B}_{2u})$	2263		2185	2141	2175
$\nu_9(\text{B}_{1u})$	2230		2145	2070	2102
$\nu_2(\text{A}_g)$					1232
$\nu_{16}(\text{B}_{2u})$	1155	1155	1180	1207	
$\nu_{10}(\text{B}_{1u})$	958				

^a Ref. 10.^b This work.^c Same wavenumbers for TCNE⁻/TBA⁺, TCNE⁻/TEA⁺, and TCNE⁻/Li⁺ systems.^d Same wavenumbers for TCNE²⁻/TBA⁺ and TCNE²⁻/TEA⁺ systems.

The C≡N str bands of TCNQ⁻ (Fig. 8) show similar behavior to those of TCNE²⁻; the wavenumbers are lower than those of TCNQ⁻ and depend upon the electrolyte. With TBAF and TEAP these bands are at 2153 and 2106 cm⁻¹, with NaClO₄ they are at 2156 and 2110 cm⁻¹, and with LiClO₄ they are at 2166 and 2118 cm⁻¹. (The positions of other bands are listed in Table 2.)

Experiments where the concentration of LiClO₄ was varied between 0.001 M and 0.1 M (while that of TBAF was half constant at 0.1 M) produced no new bands, suggesting that a complex of only one stoichiometry, probably 1:1, is formed.

The same experiments were carried out in DMSO with TBAF and LiClO₄ as electrolytes. In this solvent no evidence for ion pairing was obtained; the half wave potentials of *both* reduction waves of TCNQ were independent of electrolyte, and the C≡N str wavenumbers of the anions TCNQ⁻ and TCNQ²⁻ did not depend on

TABLE 2

Infrared vibrational wavenumbers of TCNQ and its anions in acetonitrile

Mode	Vibrational wavenumbers/cm ⁻¹				
	TCNQ	TCNQ ⁻ ^a	TCNQ ²⁻ /TBA ⁺ ^b	TCNQ ²⁻ /Na ⁺	TCNQ ²⁻ /Li ⁺
ν_{19-u}	2224	2182	2153	2156	2166
$\nu_{33}(\text{B}_{2u})$	2224	2156	2106	2110	2118
			1576		
	1545	1506	1501		1501
			1493		1493
		1362	1296		1296
		1209	1225		1235
		986			
	845	833	821		821

^a Same wavenumbers for TCNQ⁻/TBA⁺, TCNQ⁻/TEA⁺, TCNQ⁻/Na⁺ and TCNQ⁻/Li⁺ systems.^b Same wavenumbers for TCNQ²⁻/TBA⁺ and TCNQ²⁻/TEA⁺ systems.

$\nu_{19}(\text{B}_{1u})$ }
 →

electrolyte either. In fact these wavenumbers (2180 and 2153 cm^{-1} for the anion, 2151 and 2104 cm^{-1} for the dianion) are the same as those of uncomplexed $\text{TCNQ}^{\cdot-}$ and TCNQ^{2-} in acetonitrile.

DISCUSSION

TCNE is planar and belongs to the D_{2h} point group. It has four $\text{C}\equiv\text{N}$ groups and therefore four $\text{C}\equiv\text{N}$ stretching modes. Two of these, the $\nu_9(\text{B}_{1u})$ and $\nu_{15}(\text{B}_{2u})$ modes, are infrared active and Raman inactive, while the other two, the $\nu_1(\text{A}_g)$ and $\nu_{19}(\text{B}_{3g})$ modes, are Raman active and infrared inactive. Similarly there are four modes of mostly C-C stretching character: IR active $\nu_{10}(\text{B}_{1u})$ and $\nu_{16}(\text{B}_{2u})$ modes, and Raman active $\nu_3(\text{A}_g)$ and $\nu_{21}(\text{B}_{3g})$ modes. The only other in-plane stretching mode is the Raman active $\nu_2(\text{A}_g)$ mode, which is mostly stretching of the central $\text{C}=\text{C}$ bond. The transmission IR spectrum [6] of neutral TCNE has bands at 2263, 2230, 1155, and 958 cm^{-1} , and these have been assigned to the ν_{15} , ν_9 , ν_{16} , and ν_{10} modes respectively. (Actually, for neither neutral TCNE nor for its anions is it really known which $\text{C}\equiv\text{N}$ stretch band corresponds to ν_{15} and which to ν_9 .) TCNE also has bands at lower wavenumbers which have been assigned to bending vibrations. If $\text{TCNE}^{\cdot-}$ and TCNE^{2-} are planar and undistorted (i.e., of D_{2h} symmetry), they should also have two IR active $\text{C}\equiv\text{N}$ stretch modes (ν_{15} and ν_9) and two IR active C-C stretch modes (ν_{16} and ν_{10}).

While there have been other assignments of the $\text{TCNE}^{\cdot-}$ bands [16], it appears simplest to assign them to the normally infrared active modes of a species with D_{2h} symmetry. Then the 2187 and 2146 cm^{-1} bands correspond to the ν_{15} and ν_9 $\text{C}\equiv\text{N}$ stretching modes, and the 1155 and 1180 cm^{-1} bands to the ν_{16} C-C stretching mode. These assignments are summarized in Table 1. The ν_{10} C-C stretch could not be observed because it is below the 1000 cm^{-1} cutoff of the CaF_2 window used here. The spectra therefore show that addition of an electron to TCNE weakens the $\text{C}\equiv\text{N}$ bonds and strengthens the C-C bonds. This is consistent with MO calculations [19-25] which show that both additional electrons go into an orbital which is antibonding w.r.t. the $\text{C}\equiv\text{N}$ bonds, bonding w.r.t. and C-C bonds, and antibonding w.r.t. the central $\text{C}=\text{C}$ bond. The spectrum of $\text{TCNE}^{\cdot-}$ in solution (Table 1) is quite different from the spectrum of $\text{TCNE}^{\cdot-}$ in solid NaTCNE [7] state. In particular, the $\text{C}\equiv\text{N}$ str wavenumbers are higher in the solid state spectrum (2222 and 2188 cm^{-1}), and the solid-state spectrum has additional bands at ca. 1400 cm^{-1} . The appearance of the bands around 1400 cm^{-1} has been attributed to infrared activation of the A_g $\text{C}=\text{C}$ stretching mode by an interaction of the anion and cation [4-7]. This interaction could also increase the $\text{C}\equiv\text{N}$ str wavenumbers in the same manner as does the interaction of Li^+ with the dianion (cf. Fig. 3). The situation is complicated, however, by the probability of bonding between adjacent $\text{TCNE}^{\cdot-}$ units in the solid state [8-10], which may also cause changes in bond orders and in selection rules.

Again, it is simplest to assign the observed bands for TCNE^{2-} to the ν_{15} , ν_9 , and ν_{16} modes, assuming that TCNE^{2-} has D_{2h} symmetry (see Table 1). This would

$\nu_1 \text{A}_g$
 $\nu_{19} \text{B}_{3g}$

$\nu_3 \text{A}_g$

A_g

$\nu_2 \text{A}_g$

$\nu_{21} \text{B}_{3g}$

A_g

mean that, as predicted by the MO calculations, addition of an electron to TCNE^{•-} causes further weakening of the C≡N bonds and further strengthening of the C-C bonds. With LiClO₄ as electrolyte, the dianion bands are at 2175, 2102, and 1232 cm⁻¹. The C≡N bands are shifted to lower wavenumber compared to the anion radical, but not by as large an amount as in the presence of TBAF. This difference is attributed to strong ion pairing of TCNE²⁻ with Li⁺ and not with (C₄H₉)₄N⁺. Presumably the Li⁺ ion either undergoes a charge-transfer interaction with TCNE²⁻ which removes some electron density from the antibonding MO, or it causes a redistribution of π -electrons within the dianion such that the C≡N bonds are strengthened. It seems likely that only one Li⁺ ion is involved in this strong interaction with TCNE²⁻, because experiments using mixtures of TBAF (0.1 M) and LiClO₄ (0.001 M to 0.1 M) showed only the two sets of C≡N stretch bands of Fig. 3. No additional bands appeared at low Li⁺ concentrations, as might be expected if there were complexes of more than one stoichiometry. (This does not rule out weak interactions of other Li⁺ ions with the complex.) The 1232 cm⁻¹ band of the complex presents a problem. It is at higher wavenumbers than the ν_{16} C-C stretch of uncomplexed TCNE²⁻, yet it is difficult to conceive of a mechanism which could simultaneously strengthen both the C≡N and C-C bonds of example TCNE²⁻. It is more likely that the 1232 cm⁻¹ band is the normally infrared inactive $\nu_2(A_g)$ C=C stretching vibration. This mode could be made infrared active by distortion of the π -electron cloud of the dianion in the direction of the cation, which is probably sitting above the molecular plane, causing vibration of the C=C bond to give rise to an oscillating dipole normal to the molecular plane. It has been proposed, in fact, that this mechanism of activation of the ν_2 mode is responsible for bands in the solid state infrared spectra of M⁺TCNE^{•-} and (Na⁺)₂TCNE²⁻ at ca. 1400 cm⁻¹ [4-6] and 1260 cm⁻¹ [7] respectively. Possible structures of the TCNE²⁻-Li⁺ complex will be considered in more detail below.

The C≡N stretching wavenumbers of solid Na₂TCNE are, quite reasonably, intermediate between those of uncomplexed TCNE²⁻ and the TCNE²⁻-Li⁺ ion pair. The dianion appears to have an IR active C=C stretching mode and an IR inactive C-C stretching mode both in the ion pair and in the solid state, which suggests that the two environments are quite similar.

TCNQ is similar to TCNE in that it is planar and of D_{2h} symmetry. Therefore TCNQ and its anions, if they are also planar, each have two infrared active C≡N stretch modes, which are labeled $\nu_{19}(B_{1u})$ and $\nu_{33}(B_{2u})$. For neutral TCNQ these two modes are coincident at 2228 cm⁻¹ [11].

There are a number of bands at lower wavenumbers than the CN str bands, and these are listed in Table 2. Because of the complexity of the molecule, and because of uncertainty in the literature [11,26], no attempt is made here to assign these other bands. The positions of all bands of TCNQ^{•-} are independent of electrolyte. The lowest unoccupied molecular orbital of TCNQ, into which the additional electrons must go, is antibonding with respect to the C≡N bonds, just as it is for TCNE. In accordance with this, the C≡N str wavenumber of TCNQ^{•-} are lower than that of TCNQ.

TABLE 3

Electrochemical and infrared data for the reductions of TCNE and TCNE^{2-} in acetonitrile

Reaction electrolyte cation	$\text{TCNE} + e^- \rightarrow \text{TCNE}^-$			$\text{TCNE}^{2-} \rightarrow \text{TCNE}^{2-}$		
	TBA ⁺	Na ⁺	Li ⁺	TBA ⁺	Na ⁺ Li ⁺	
$E_{1/2}/V$ (vs. Ag/Ag ⁺)	-0.07	-0.07	-0.07	-1.05	-0.94	-0.79
ν_{CN} decrease/ cm^{-1}	82		82	59		26
ν_{CN} splitting/ cm^{-1}	40		40	71		73

The anticipated weakening of the C≡N bonds occurs upon adding the second electron, but the wavenumber shifts are smaller in the presence of alkali metal cations, and once again this is attributed to complex formation with the small cations.

The SNIPTIRS spectrum of TCNQ^{2-} (Table 2) agrees very well with the IR spectrum of LiTCNQ in acetonitrile [26]. The solid state IR spectra of alkali metal salts of TCNQ^{2-} [11,26] show C≡N stretching wavenumbers which are higher than those of solution free TCNQ^{2-} , and which increase with decreasing size of the cation, presumably because of interactions between the cations and the anion.

There are close similarities between the solution spectra of TCNQ^{2-} (Table 2) and the solid state spectrum of Na_2TCNQ [11], which has strong bands at 2164, 2096, 1503, 1498, 1303, 1238, and 822 cm^{-1} , and weaker bands at 1598, 1435, and 1353 cm^{-1} . The average C≡N str wavenumber of Na_2TCNE (2130 cm^{-1}) is close to that of the $\text{TCNQ}^{2-}\text{-Na}^+$ ion pair (2133 cm^{-1}), but the splitting is greater in the solid state.

Cyclic voltammetry provided the first evidence for ion pairing between alkali metal cations and the dianions of TCNE and TCNQ in acetonitrile; the half wave potentials ($E_{1/2}$) of the second reduction waves varied with electrolyte, while the half wave potentials of the first reduction waves did not [3]. Further evidence for ion pairing comes from the electrolyte dependence of the C≡N stretching wavenumbers of TCNE^{2-} and TCNQ^{2-} . Tables 3-5 compare the half wave potential of each reduction wave in the TCNE/AN, TCNQ/AN, and TCNQ/DMSO systems with the difference in average C≡N str wavenumber between the oxidized and reduced species. Also shown is the splitting between the two C≡N str bands of the reduced species. A number of points are apparent from these Tables:

TABLE 4

Electrochemical and infrared data for the reductions of TCNQ and TCNQ^{2-} in acetonitrile

Reaction electrolyte cation	$\text{TCBQ} + e^- \rightarrow \text{TCNQ}^{2-}$			$\text{TCNQ}^{2-} + e^- \rightarrow \text{TCNQ}^{2-}$		
	TBA ⁺	Na ⁺	Li ⁺	TBA ⁺	Na ⁺	Li ⁺
$E_{1/2}/V$ (vs. Ag/Ag ⁺)	-0.11	-0.11	-0.11	-0.66	-0.63	-0.52
ν_{CN} decrease/ cm^{-1}	55	55	55	39	36	27
ν_{CN} splitting/ cm^{-1}	26	26	26	47	47	48

TABLE 5

Electrochemical and infrared data for the reductions of TCNQ and TCNQ^{•-} in DMSO


Reaction electrolyte cation	TCNQ + e ⁻ → TCNQ ^{•-}		TCNQ ^{•-} + e ⁻ → TCNQ ²⁻	
	TBA ⁺	Li ⁺	TBA ⁺	Li ⁺
$E_{1/2}/V$ vs. SCD	+0.29	+0.29	-0.26	-0.26
ν_{CN} decrease/cm ⁻¹	54	54	39	39
ν_{CN} splitting/cm ⁻¹	27	27	47	47

(i) For neither anion radical is there any evidence of ion pairing with alkali metal cations.


(ii) The half wave potentials for reduction of the anion radicals to the dianions [3] in acetonitrile become less negative when TBAF or TEAP electrolytes are replaced with lithium or sodium perchlorate. This shows that the dianions are stabilized by interaction with the small alkali metal cations.

(iii) Li⁺ causes a greater positive shift in $E_{1/2}$ than does Na⁺ [3]. It was determined that the ion pairs are of the contact rather than the solvent separated type.

(iv) When the half wave potentials indicate that ion pairs are formed, shifts in the C≡N str wavenumbers are also observed. In fact, ion pairing causes a shift to higher wavenumber which is roughly proportional to the positive shift in $E_{1/2}$. To see why this is so, we point out pertinent information regarding the structures of the ion pairs. It is most likely that only one cation interacts strongly with each dianion, because decreasing the the concentration of metal ion fails to reveal any new bands. This cation would probably be situated in the center of the π -electron cloud above the molecular plane of the dianion; a less symmetrical location would give rise to four IR active C≡N stretch modes, instead of the two which are observed. (This location would also nicely account for the infrared activation of the central C=C bond in the TCNE²⁻-Li⁺ complex.) According to a simple electrostatic model, placing a cation over the central C=C bond of TCNE²⁻ would increase the contribution of resonance structures such as



relative to structures such as



Similar considerations apply to TCNQ^{2-} ; placement of a cation over the center of the six-membered ring would increase contributions from resonance structures where the $\text{C}\equiv\text{N}$ triple bond is intact. This accounts qualitatively for the increase in $\text{C}\equiv\text{N}$ str wavenumber caused by ion pairing. (Note that if the cation was associated with an N atom, a decrease in $\text{C}\equiv\text{N}$ str wavenumber would be expected.) According to this model, then, the shift in $E_{1/2}$ and shift in $\text{C}\equiv\text{N}$ str wavenumber are related; neglecting changes in solvation of the dianion and cation, the shift in $E_{1/2}$ measures the energy of electrostatic interaction of the cation with the negative charges on the dianion, and the shift in $\bar{\nu}_{\text{CN}}$ measures the ability of the cation to pull negative charge off the nitrogen atoms and towards the center of the molecule. Therefore both the shift in $E_{1/2}$ and the shift in $\bar{\nu}_{\text{CN}}$ increase with decreasing size of the cation and decreasing size of the dianion.

(v) While the average position of the $\text{C}\equiv\text{N}$ str bands of the dianions are affected by ion pairing, their splitting is predicted to be independent of ion pairing interactions. In fact it appears that the difference in the two $\text{C}\equiv\text{N}$ stretch bands for the dianions is directly proportional to the charge per π -bonded atom in the molecule, the constant of proportionality being 390 cm^{-1} .

(vi) There is no strong ion pairing between TCNQ^{2-} and Li^+ in DMSO. Presumably the loss of solvation of Li^+ upon ion pair formation makes this process unfavorable in DMSO. (The solvation energy of Li^+ is higher in DMSO than in acetonitrile, the Gutman donor numbers in these two solvents being 30 and 14, respectively [27].)

CONCLUSIONS

(i) Strong contact ion pairing takes place in acetonitrile between alkali metal cations and the dianions of TCNE and TCNQ. The spectra of the ion pairs are quite similar to the solid state spectra of the dianions.

(ii) Ion pairing shifts the $\text{C}\equiv\text{N}$ stretch bands of TCNE^{2-} and TCNQ^{2-} to higher wavenumber without changing the number of these bands nor the splitting between them. This is consistent with location of a cation above the central $\text{C}=\text{C}$ bond in TCNE^{2-} and above the six-membered ring in TCNQ^{2-} .

(iii) Ion pairing appears to make infrared active the (totally symmetric) $\text{C}=\text{C}$ stretching vibration of TCNE^{2-} .

(iv) The spectrum of TCNE^{2-} in acetonitrile is the same in the presence of TEAP as it is in the presence of TBAF. This means that the large difference in the standard heterogeneous rate constant of the $\text{TCNE}^{\cdot-}\text{TCNE}^{2-}$ couple in these two systems [3] cannot be due to differences in the structure of the dianion, and must instead be due to surface effects such as distance dependent electron transfer rates or different potential distributions in the double layer.

(v) There is no strong ion pairing in acetonitrile or DMSO between alkali metal cations and the anion radicals of TCNE and TCNQ. The spectra are very different from the solid state spectra of $\text{TCNE}^{\cdot-}$ and $\text{TCNQ}^{\cdot-}$ for that reason.

(vi) There is no strong ion pairing in DMSO between Li^+ and TCNQ^{2-} , probably because of strong solvation of Li^+ by DMSO.

REFERENCES

- 1 (a) S. Pons, J.K. Foley, J. Russell and M. Severson, in J.O'M. Bockris and B.G. Conway (Eds.), *Modern Advances in Electrochemistry*, Plenum Press, New York, 1968, Ch. 3; (b) J.K. Foley, C. Korzeniewski, J.L. Daschbach and S. Pons, in A.J. Bard (Ed.), *Electroanalytical Chemistry*, Vol. 14, Marcel Dekker, New York, 1986, Ch. 4.
- 2 J. Ashley, J.K. Foley, S.B. Khoo and S. Pons, unpublished work.
- 3 (a) S.B. Khoo, Ph.D. Thesis, University of Alberta, 1984; (b) S.B. Khoo, J.K. Foley and S. Pons, *J. Electroanal. Chem.*, 215 (1986) 273.
- 4 J. Stanley, D. Smith, B. Latimer and J.P. Devlin, *J. Phys. Chem.*, 70 (1966) 2011.
- 5 J.C. Moore, D. Smith, Y. Youhne and J.P. Devlin, *J. Phys. Chem.*, 75 (1971) 325.
- 6 J.J. Hinkel and J.P. Devlin, *J. Chem. Phys.*, 58 (1973) 4750.
- 7 M.S. Khatkale and J.P. Devlin, *J. Phys. Chem.*, 83 (1979) 1636.
- 8 M. Itoh, *Bull. Chem. Soc. Jpn.*, 45 (1972) 1947.
- 9 K. Yokoyama, S. Maeda, C. Etoh, S. Matsuzaki and K. Toyoda, *Bull. Chem. Soc. Jpn.*, 53 (1980) 36.
- 10 S. Matsuzaki, T. Mitsuishi and K. Toyoda, *Chem. Phys. Lett.*, 91 (1982) 296.
- 11 M.S. Khatkale and J.P. Devlin, *J. Chem. Phys.*, 70 (1979) 1851.
- 12 D.H. Eargle and R. Emrich, *J. Org. Chem.*, 35 (1970) 3744.
- 13 D.L. Jeanmaire, M.R. Suchanski and R.P. Van Duyne, *J. Am. Chem. Soc.*, 97 (1975) 1699.
- 14 D.L. Jeanmaire and R.P. Van Duyne, *J. Am. Chem. Soc.*, 98 (1976) 4029.
- 15 R.P. Van Duyne, M.R. Suchanski, J.M. Lakovita, A.R. Siedle, K.D. (Parks and T.M. Cotton, *J. Am. Chem. Soc.*, 101 (1979) 2832.
- 16 S. Pons, S.B. Khoo, A. Bewick, M. Datta, J.J. Smith and G. Zachman, *J. Phys. Chem.*, 88 (1984) 3575.
- 17 M. Datta, R.W. Jasson and J.J. Freeman, *Spectrosc. Lett.*, 19 (1986) 129.
- 18 H. Lund and P. Iverson, in M. Baizer (Ed.), *Organic Electrochemistry*, Marcel Dekker, New York, 1969, p. 57.
- 19 P.H. Rieger and G.K. Fraenkel, *J. Chem. Phys.*, 37 (1962) 2795.
- 20 Y. Iida, *Bull. Chem. Soc. Jpn.*, 46 (1973) 423.
- 21 B.R. Penfold and W.N. Lipscomb, *Acta Crystallogr.*, 14 (1961) 589.
- 22 R.C. Haddon, *Aust. J. Chem.*, 28 (1975) 2333.
- 23 J. Ladik, A. Karpfen, G. Stoffhoff and P. Fulde, *Chem. Phys.*, 7 (1975) 267.
- 24 M.J.S. Dewar and H.S. Rzepa, *J. Am. Chem. Soc.*, 100 (1978) 784.
- 25 K. Korch-Jespersen and M.A. Ratner, *Theor. Chim. Acta*, 47 (1978) 283.
- 26 R. Bozio, A. Girlando and C. Pecile, *J. Chem. Soc., Faraday Trans. II*, 71 (1975) 1237.
- 27 V. Gutmann, *Chimia*, 31 (1977) 1.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 92132	1

ABSTRACTS DISTRIBUTION LIST, SDIO/IST

Dr. Robert A. Osteryoung
Department of Chemistry
State University of New York
Buffalo, NY 14214

Dr. Douglas N. Bennion
Department of Chemical Engineering
Brigham Young University
Provo, UT 84602

Dr. Stanley Pons
~~Department of Chemistry~~
~~University of Utah~~
Salt Lake City, UT 84112

Dr. H. V. Venkatesetty
Honeywell, Inc.
10701 Lyndale Avenue South
Bloomington, MN 55420

Dr. J. Foos
EIC Labs Inc.
111 Downey St.
Norwood, MA 02062

Dr. Neill Weber
Ceramatec, Inc.
163 West 1700 South
Salt Lake City, UT 84115

Dr. Subhash C. Narang
SRI International
333 Ravenswood Ave.
Menlo Park, CA 94025

Dr. J. Paul Pemsler
Castle Technology Corporation
52 Dragon Ct.
Woburn, MA 01801

Dr. R. David Rauh
EIC Laboratory Inc.
111 Downey Street
Norwood, MA 02062

Dr. Joseph S. Foos
EIC Laboratories, Inc.
111 Downey Street
Norwood, Massachusetts 02062

Dr. Donald M. Schleich
Department of Chemistry
Polytechnic Institute of New York
333 Jay Street
Brooklyn, New York 01

Dr. Stan Szpak
Code 633
Naval Ocean Systems Center
San Diego, CA 92152-5000

Dr. George Blomgren
Battery Products Division
Union Carbide Corporation
25225 Detroit Rd.
Westlake, OH 44145

Dr. Ernest Yeager
Case Center for Electrochemical
Science
Case Western Reserve University
Cleveland, OH 44106

Dr. Mel Miles
Code 3852
Naval Weapons Center
China Lake, CA 93555

Dr. Ashok V. Joshi
Ceramatec, Inc.
2425 South 900 West
Salt Lake City, Utah 84119

Dr. W. Anderson
Department of Electrical &
Computer Engineering
SUNY - Buffalo
Amherst, Massachusetts 14260

Dr. M. L. Gopikanth
Chemtech Systems, Inc.
P.O. Box 1067
Burlington, MA 01803

Dr. H. F. Gibbard
Power Conversion, Inc.
495 Boulevard
Elmwood Park, New Jersey 07407

DL/1113/87/2

ABSTRACTS DISTRIBUTION LIST, SDIO/IST

Dr. V. R. Koch
Covalent Associates
52 Dragon Court
Woburn, MA 01801

Dr. Randall B. Olsen
Chronos Research Laboratories, Inc.
4186 Sorrento Valley Blvd.
Suite H
San Diego, CA 92121

Dr. Alan Hooper
Applied Electrochemistry Centre
Harwell Laboratory
Oxfordshire, OX11 0RA UK

Dr. John S. Wilkes
Department of the Air Force
The Frank J. Seiler Research Lab.
United States Air Force Academy
Colorado Springs, CO 80840-6528

Dr. Gary Bullard
Pinnacle Research Institute, Inc.
10432 N. Tantan Avenue
Cupertino, CA 95014

Dr. J. O'M. Bockris
Ementech, Inc.
Route 5, Box 946
College Station, TX 77840

Dr. Michael Binder
Electrochemical Research Branch
Power Sources Division
U.S. Army Laboratory Command
Fort Monmouth, New Jersey 07703-5000

Professor Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton, Hants, SO9 5NH UK